

# Patient Education - Minoxidil 3mg/Dutasteride 0.1mg Capsule

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Only use medication as prescribed

This medication was compounded by a pharmacy per prescriber request. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain.

Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss a dose, take as soon as you remember. If it is close to the next dose, skip the missed dose. Do not take double the dosage.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program - Website: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Please review the following:

Minoxidil - is used to help you regrow hair on your scalp & may also be used for purposes not listed in this medication guide. Minoxidil will not cause permanent regrowth of scalp hair. You must continue using the product to keep the regrowth of your hair.

Using more of this medicine than recommended will not speed up hair growth, and may cause dangerous side effects. Ask a doctor or pharmacist if it is safe for you to use minoxidil if you have other medical conditions, especially heart disease.

FDA pregnancy category C. It is not known whether minoxidil topical will harm an unborn baby. Do not use this medicine without a doctor's advice if you are pregnant. It is not known whether minoxidil passes into breast milk or if it could harm a nursing baby. Do not use this medicine without a doctor's advice if you are breast-feeding a baby.

Do not use this medicine on anyone under 18 years old without medical advice.

Call your doctor at once if you have: Chest pain, quick drop of blood pressure fast heartbeats, swelling in your hands or feet, rapid weight gain, a light-headed feeling, like you might pass out headache, dizziness, confusion, or flushing (warmth, redness, or tingly feeling).

Common side effects may include: hair growth (possibly in various parts of the body), lowered blood pressure

Dutasteride- A 5 Alpha-Reductase Inhibitor, used to help you regrow hair on your scalp by preventing the conversion of testosterone to dihydrotestosterone (DHT) in the body. Dutasteride may also be used for purposes not listed in this medication guide.

Dutasteride can cause birth defects if a woman is exposed to it during pregnancy & should not be handled by a woman who is pregnant or may become pregnant. If a woman accidentally comes into contact with Dutasteride wash the area with soap and water right away.

Dutasteride should never be taken by a child. If a child accidentally comes into contact with Dutasteride wash the area with soap and water right away.

You should not use Dutasteride if you are allergic to it.

Using Dutasteride may increase your risk of developing prostate cancer in males. Talk with your doctor if you have concerns about this risk. Tell your doctor if you have ever had: liver disease, or abnormal liver function tests, prostate cancer, a bladder muscle disorder, stricture of your urethra urination problems.

Get emergency medical help if you have signs of an allergic reaction: hives difficulty breathing swelling of your face, lips, tongue, or throat.

Call your doctor at once if you notice any signs of male breast cancer, such as: breast lumps, breast pain or tenderness, nipple discharge or any other breast changes.

Common side effects may include: loss of interest in sex impotence trouble having an orgasm or abnormal ejaculation.

The sexual side effects of Dutasteride may continue after you stop taking Dutasteride. Talk to your doctor if you have concerns about these side effects. This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.