

# Patient Education - Sulfacetamide Sodium 10%, Tacrolimus 0.1% Topical Gel

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

## Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes. Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program

Website address: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Sulfacetamide Sodium - an antibiotic that fights bacteria on the skin.

Sulfacetamide sodium topical (for the skin) is used to treat acne, dandruff, seborrheic dermatitis (red, flaking skin rash), and certain skin infections.

Sulfacetamide sodium topical may also be used for purposes not listed in this medication guide.

You should not use this medicine if you are allergic to sulfacetamide sodium or sulfa drugs.

To make sure this medicine is safe for you, tell your doctor if you have: lupus asthma or a sulfite allergy.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether sulfacetamide sodium topical passes into breast milk or if it could harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Do not use this medicine on a child younger than 12 years old without direction of a prescriber.

Wash your hands after applying this medicine.

Avoid getting this medicine on your clothing. Sulfacetamide sodium topical may leave a slight yellow color on light fabrics.

Avoid exposure to sunlight or tanning beds. Sulfacetamide sodium topical can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Avoid using other medications on the areas you treat with sulfacetamide sodium topical unless your doctor tells you to.

Stop using this medicine and call your doctor at once if you have:

the first sign of any skin rash, no matter how mild pale skin, easy bruising or bleeding new or worsening skin infection

liver problemsnausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

lupus-like syndromejoint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color or

severe skin reactionfever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

Common side effects may include:

mild swelling, itching, or redness of treated skin dry skin or yellow discoloration of treated skin (may be a sign that you have applied too much of the medicine).

Tacrolimus - an immunosuppressant. It works by decreasing your body's immune system & is used to treat severe atopic dermatitis (eczema).

Tacrolimus may also be used for purposes not listed in this medication guide.

#### Important Information

You should not use tacrolimus topical if you are allergic to it.

Before using tacrolimus topical, tell your doctor if you have skin cancer or a skin infection (including herpes or chickenpox), any genetic skin disorder (such as Netherton's syndrome), a weak immune system, kidney disease, or swelling, redness, or irritation of large areas of your skin.

Tacrolimus can lower the blood cells that help your body fight infections. This can make it easier for you to get sick from a virus such as chicken pox or herpes (cold sores or shingles). Tell your doctor if you have been exposed to any illness.

Avoid sunlight, sun lamps, tanning beds, and phototherapy treatments with UVA or UVB light. If you must be outdoors, wear loose clothing over the skin areas treated with tacrolimus topical. Do not use sunscreen on treated skin unless your doctor has told you to.

Stop using tacrolimus and call your doctor at once if you have a serious side effect such as:

severe stinging, burning, itching, or soreness where the medicine is applied swollen glands redness or crusting around your hair follicles or signs of a skin infection (redness, swelling, itching, oozing).

Less serious side effects may include:

mild burning, stinging, or itching skin redness acne cold or flu symptoms such as stuffy nose, sneezing, sore throat headache or feeling more sensitive to hot or cold temperatures.