

# Patient Education - Tacrolimus 0.09%, Dapsone 6.5%, Calcipotriene 0.005%, Topical Ointment

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

- Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes. Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program

Website address: <https://www.fda.gov/Safety/MedWatch/default.htm>

Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Tacrolimus - an immunosuppressant. It works by decreasing your body's immune system & is used to treat severe atopic dermatitis (eczema).

Tacrolimus may also be used for purposes not listed in this medication guide.

## Important Information

You should not use tacrolimus topical if you are allergic to it.

Before using tacrolimus topical, tell your doctor if you have skin cancer or a skin infection (including herpes or chickenpox), any genetic skin disorder (such as Netherton's syndrome), a weak immune system, kidney disease, or swelling, redness, or irritation of large areas of your skin.

Tacrolimus can lower the blood cells that help your body fight infections. This can make it easier for you to get sick from a virus such as chicken pox or herpes (cold sores or shingles). Tell your doctor if you have been exposed to any illness.

Avoid sunlight, sun lamps, tanning beds, and phototherapy treatments with UVA or UVB light. If you must be outdoors, wear loose clothing over the skin areas treated with tacrolimus topical. Do not use sunscreen on treated skin unless your doctor has told you to.

Stop using tacrolimus and call your doctor at once if you have a serious side effect such as:

severe stinging, burning, itching, or soreness where the medicine is applied swollen glands redness or crusting around your hair follicles or signs of a skin infection (redness, swelling, itching, oozing).

Less serious side effects may include:

mild burning, stinging, or itching skin redness acne cold or flu symptoms such as stuffy nose, sneezing, sore throat headache or feeling more sensitive to hot or cold temperatures.

Dapsone - an anti-infective medication.

Dapsone topical (for the skin) is used to treat acne & may also be used for purposes not listed in this medication guide.

To make sure dapsone topical is safe for you, tell your doctor if you have:

glucose 6-phosphate dehydrogenase (G6PD) deficiency a blood disorder called methemoglobinemia (abnormal red blood cells in your blood) or if you have ever had methemoglobinemia in the past.

It is not known whether dapson topical will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant. Dapson topical can pass into breast milk and may harm a nursing baby. You should not breast-feed while using this medicine.

Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat.

Dapson topical may cause methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a), a serious condition in which the amount of oxygen in your blood stream becomes dangerously low. Stop using this medicine and call your doctor at once if you notice a blue or gray appearance of your lips, fingernails, or the inside of your mouth.

Call your doctor at once if you have: Severe sore throat unusual thoughts or behavior a red blood cell disorderback pain, pale or yellowed skin, dark colored urine, fever, confusion or weakness pancreas problemssevere pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate or

severe skin reactionfever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

Common side effects may include: Dry or peeling skin oily skin or redness where the medicine was applied.

Calcipotriene - Calcipotriene is a man-made form of vitamin D. It is involved in the growth and development of skin cells.

It is used to treat plaque psoriasis (psoriasis with scaly patches) and may also be used for purposes not listed in this medication guide.

Do not take by mouth. Calcipotriene topical is for use only on the skin. Do not use calcipotriene topical on open wounds or on sunburned, windburned, dry, chapped, or irritated skin.

Tell your doctor if you are pregnant or breastfeeding.

Do not use this medicine on a child without medical advice.

Along with its needed effects, calcipotriene topical may cause some unwanted effects. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects.

Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat.

Stop using calcipotriene topical and call your doctor at once if you have:

-High calcium levelsconfusion, tiredness, nausea, vomiting, loss of appetite, constipation, increased thirst or urination, weight loss.

Common side effects may include:

-Mild skin irritation skin rash or itching.

Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

-Burning, dryness, irritation, peeling, or redness of the skin

-Pain at the application site