

Patient Education - Cidofovir 2% Topical Ointment

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 - Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes.

Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The "Do not use after" date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program - Website: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Cidofovir is an antiviral medicine that is used to treat infections caused by viruses, like cold sores and may also be used for purposes not listed in this medication guide.

You should not be treated with cidofovir if you are allergic to it, or if you have: moderate to severe kidney disease or a history of severe allergic reaction to probenecid or sulfa drugs.

Tell your doctor if you have ever had: kidney disease diabetes or pancreas problems.

If you are a woman, do not use cidofovir if you are pregnant. Use effective birth control to prevent pregnancy while you are using this medicine and for at least 1 month after your last dose.

Cidofovir side effects: Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have: vision changes, white or yellow patches on your eyes kidney problems swelling, rapid weight gain, little or no urinating, increased thirst, loss of appetite, constipation, pain in your side or lower back low blood cell counts fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath or pancreatitis severe pain in your upper stomach spreading to your back, nausea and vomiting.

Common side effects may include: nausea, vomiting, diarrhea pain, weakness rash headache or hair loss.

You may have side effects such as: nausea, vomiting skin rash fever, chills or headache.

This is not a complete list of side effects and others may occur.